

RATES FOR NON-PERTURBATIVE PROCESSES IN HOT NON-ABELIAN PLASMAS^a

PETER ARNOLD

Department of Physics, University of Virginia, Charlottesville, VA 22901

overviewing work done¹ in collaboration with

DAM SON

*Center for Theoretical Physics, Department of Physics,
Massachusetts Institute of Technology, Cambridge, MA 02139*

LAURENCE G. YAFFE

Department of Physics, University of Washington, Seattle, Washington 98195

I give a simple physical picture of why the rate Γ per unit volume of non-perturbative processes in hot non-Abelian plasmas (such as electroweak baryon number violation in the very early universe) depends on coupling as $\Gamma \sim \alpha^5 T^4 \ln(1/\alpha)$. The logarithmic dependence was recently discovered by Bödeker.

1 Introduction

I will be discussing the rate for non-perturbative processes in hot non-Abelian plasmas. For me “hot” means high enough temperature that the running coupling $\alpha(T)$ is small, that chemical potentials are ignorable, and that there is no spontaneous symmetry breaking. Examples of non-perturbative processes include chirality violation in hot QCD and baryon number violation in hot electroweak theory (in its high-temperature symmetric phase). Both cases are related by the anomaly

$$\Delta Q \sim g^2 \int d^4x \operatorname{tr} F \tilde{F} \quad (1)$$

to non-perturbative fluctuations of the gauge fields. The question then becomes: what is the rate for such non-perturbative fluctuations?

In fact, it turns out that a non-trivial question is how the rate Γ per unit volume even depends on the coupling α for small coupling:

$$\Gamma = \# \alpha^? T^4 \quad (+ \text{ higher-order}). \quad (2)$$

The T^4 dependence just comes from dimensional analysis. In this talk, I shall review the spatial and temporal scales characteristic of non-perturbative

^aTalk given at “Continuous Advances in QCD ’98.”

physics in the hot plasma, showing that the distance scale is

$$R \sim \frac{1}{g^2 T} \quad (3)$$

and the time scale is

$$t \sim \frac{1}{g^4 T \ln(1/g)}. \quad (4)$$

If you accept these scales on faith for a moment, then one will have, roughly, one unsuppressed non-perturbative process per volume R^3 per time t , so that

$$\Gamma \sim \frac{1}{R^3 t} \sim \alpha^5 T^4 \ln\left(\frac{1}{\alpha}\right). \quad (5)$$

The α^5 is a result of my collaborators and I from two years ago.² The logarithmic enhancement is a recent and interesting result of Dietrich Bödeker.³

2 The Spatial Scale

Imagine a non-perturbative fluctuation of the gauge fields with some spatial extent R . Non-perturbative means that gA is not a perturbation in the covariant derivative $D = \partial - gA$ and so

$$A \sim \frac{1}{gR}. \quad (6)$$

The energy of the configuration is then order

$$E \sim \frac{1}{g^2 R}. \quad (7)$$

The probability of such a fluctuation at finite temperature T is, roughly speaking, given by a Maxwell-Boltzmann factor:

$$e^{-\beta E} \sim e^{-1/(g^2 RT)}, \quad (8)$$

and this probability is unsuppressed when

$$R \gtrsim \frac{1}{g^2 T}. \quad (9)$$

In fact, because of entropy, the *smallest* unsuppressed distance scale is the dominant one—there are more possible small configurations than big ones. So the characteristic scale of non-perturbative fluctuations is $R \sim 1/g^2 T$ as claimed in the introduction, corresponding to a spatial momentum scale of

$$k \sim g^2 T. \quad (10)$$

Some readers may wonder how non-perturbative physics in the hot plasma is consistent with deconfinement—something that’s supposed to happen when you heat up non-Abelian gauge theory. In a hot plasma, static electric fields are screened by the Debye effect, which arises from the response of the charges in the plasma to the electric field. This screening of the electric field is what is meant when people speak of “deconfinement”—it means that the effective potential between static test charges is not confining. However, static magnetic fields are *not* screened in a plasma (which is why, for instance, the galaxy can have a magnetic field). The physics associated with static or nearly static non-Abelian magnetic fields can still be non-perturbative.^b

3 The time scale

It was long assumed in the literature on electroweak baryogenesis that the time scale t was the same as the spatial scale R . This need not be true because the presence of the plasma breaks Lorentz invariance. In fact, the dynamics is slowed down because plasmas are conductors and because of Lenz’s Law: a conductive medium resists changes in magnetic field. This qualitative observation, in this context, is due to Guy Moore.⁴ Now to understand the time scale, start with Ampere’s Law:

$$\nabla \times \mathbf{B} = \frac{d\mathbf{E}}{dt} + \mathbf{J} = \frac{d\mathbf{E}}{dt} + \sigma\mathbf{E}, \quad (11)$$

where σ is the (color) conductivity of the plasma and where, for the sake of notational and conceptual simplicity in these rough arguments, I will simply write familiar Abelian equations (*e.g.* $\nabla \times \mathbf{B}$ instead of $\mathbf{D} \times \mathbf{B}$). Let’s anticipate the end result that the dynamics will be slow and ignore the $d\mathbf{E}/dt$ term above. In $A_0 = 0$ gauge, the result may be written as

$$\nabla \times \nabla \times \mathbf{A} \simeq -\sigma \frac{d\mathbf{A}}{dt}. \quad (12)$$

In terms of characteristic scales, this means

$$\frac{1}{R^2} A \sim \sigma \frac{1}{t} A, \quad (13)$$

and so the characteristic time scale is

$$t \sim \sigma R^2. \quad (14)$$

^b On a more technical level, large space-like Wilson loops still have area-law behavior at high temperature.




Figure 1: A spatially oscillating electric field with wave number k .

As I shall review, the color conductivity of a non-Abelian plasma is of order

$$\sigma \sim \frac{T}{\ln(1/\alpha)}. \quad (15)$$

Combined with the spatial scale (3), this leads to the results (4) and (5) claimed for the time scale and the rate Γ in the introduction. A more proper and detailed discussion of the above and of what follows may be found in ref. 5.

4 The color conductivity

4.1 Collisionless Plasmas

I'll now explain how one understands the conductivity in hot plasmas. I'll begin by first considering the case of a collisionless plasma—that is, what is the current response to an external electric field if we ignore interactions between the charge carriers in the plasma? Consider this response as a function of the wave number k . Specifically, imagine applying an external electric field of the form $\mathbf{E}(\mathbf{x}) = \mathbf{E}_0 \cos(kz)$, such as depicted in fig. 1. Charges in the plasma will be accelerated by the electric field, creating a current. If the electric field were spatially homogeneous, the charge carriers would be accelerated indefinitely, and the current would grow arbitrarily large in the absence of collision. In the case of the spatially oscillating electric field, however, this growth of the current is cut off when the particles fly into a region where the direction of $\mathbf{E}$ is reversed. The larger k , the sooner this happens, and the smaller the current. The conductivity $\sigma = J/E$ turns out to be of order

$$\sigma \sim \frac{g^2 T^2}{k}, \quad (16)$$




Figure 2: The path of a drunk heading towards a warehouse of wine.

where the $1/k$ embodies the cut-off effect just described. One factor of g is from the coupling of the charges to the electric field, the other factor of g is from the definition of the current as charge times velocity.

If one plugs the spatial scale (10) of soft dynamics into (16), the order of the conductivity becomes $\sigma \sim T$, which would give a time scale (14) of $t \sim 1/g^4 T$ and rate $\Gamma \sim \alpha^5 T^4$. This answer misses Bödeker's logarithmic enhancement.

4.2 What about $k \rightarrow 0$?

First consider the case of an Abelian plasma. The collisionless result (16) obviously makes no sense as we take $k \rightarrow 0$. Real conductors, like copper, obviously have a finite response to homogeneous electric fields. The reason is collisions. Every so often, electrons have hard, large-angle collisions that randomize their direction. Each such randomization of direction randomizes that electron's contribution to the current. This situation is depicted in fig. 2, which shows the path taken by a drunk who is being accelerated to the right by the smell of a warehouse full of wine. Every so often the drunk smashes into a lamp post and stumbles away in a random direction. As a result, the drunk's velocity towards the warehouse is limited.

Here, the time scale that determines the limiting current is the mean free time for the direction to be randomized. In the collisionless case, it was instead the time scale of order $1/k$ for particles to travel from a region of $+\mathbf{E}_0$ to $-\mathbf{E}_0$. A diagram for a collision is shown in fig. 3. The square matrix element $|\mathcal{M}|^2$ is explicitly of order g^4 . The mean free time for hard collisions is then order

$$\tau_{\text{large}} \sim \frac{1}{g^4 T \ln}, \quad (17)$$




Figure 3: A collision between charge carriers.

where the T is by dimensional analysis and the logarithm turns out to arise because randomization of direction can occur either through a single large-angle scattering or a succession of (individually more probable) small-angle ones. If this were the end of the story, then collisions would be irrelevant for non-perturbative dynamics in non-Abelian plasmas, because the collision time scale τ_{large} is large compared to the time scale $1/k \sim 1/g^2 T$ relevant to the previous section. That is, collisions would be rare enough that we could ignore them, at the spatial scale of interest, and use the collisionless result (16) for the conductivity.

However, as first noted by Selikhov and Gyulassy,^{6c} the situation is quite different for non-Abelian plasmas. The reason is that the virtual gluon exchanged in fig. 3 carries color and so changes the colors of the colliding particles. That means that very soft, small-angle collisions, even though they do not individually randomize the *direction* of the charge carriers, can instead randomize the *charge* of the charge carriers. Randomizing the charges, however, is enough to randomize the individual currents. Imagine that the drunk in fig. 2 now has a multiple personality disorder: sometimes he’s a drunk (“wine! more wine!”), sometimes he’s a teetotaler (“Goodness Gracious, I must head home”), and he vacillates between these personalities much faster than he has hard collisions with lamp posts.^d

The mean free time that determines the color conductivity is therefore the mean free time for *any* angle scatterings, not just large-angle ones. The total cross-section for any scattering suffers from the usual Coulomb divergence and would formally be infinite if not for screening effects in the plasma and non-perturbative effects. When one includes such screening effects, one finds that the mean free time is

$$\tau_{\text{any}} \sim \frac{1}{g^2 T \ln}. \quad (18)$$

^c I am grateful to H. Heiselberg for pointing me towards the literature on this subject after my talk at the conference.

^d The author is fully aware that this analogy with the drunk serves no useful purpose—he just enjoys it.

This time is *smaller* than $1/k$ by a logarithm and so dominates the physics when α is small enough that the logarithm is big. Replacing the time scale $1/k$ by τ_{any} in eq. (16) for the conductivity, one obtains

$$\sigma \sim \frac{T}{\ln(1/\alpha)} \quad (19)$$

as claimed earlier. The fact that τ_{any} beats $1/k$ by a logarithm is what, in this language, is responsible for the logarithmic enhancement of the rate Γ in (5), found by Bödeker.

For a much more in depth discussion of the topics I have discussed, see ref. 1.

References

1. P. Arnold, D. Son, and L. Yaffe, **hep-ph/9810216**, Univ. of Washington report no. UW/PT 98-10.
2. P. Arnold, D. Son, and L. Yaffe, **hep-ph/9609481**, *Phys. Rev.* **D55**, 6264 (1997).
3. D. Bödeker, **hep-ph/9810430**, *Phys. Lett.* **B426**, 351 (1998).
4. Guy Moore, private communication.
5. P. Arnold, D. Son, and L. Yaffe, University of Washington preprint UW/PT 98-10.
6. A. Selikhov and M. Gyulassy, **nucl-ph/9307007**, *Phys. Lett.* **B316**, 373 (1993).